

TITLE OF THE INVENTION

METHOD OF PRODUCING A BIOSENSOR PROTEIN CAPABLE OF
REGULATING A FLUORESCENCE PROPERTY OF GREEN FLUORESCENT
PROTEIN, AND THE BIOSENSOR PROTEIN PRODUCED BY
5 THE METHOD

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is based upon and claims the
benefit of priority from the prior Japanese Patent
Application No. 2000-356047, filed November 22, 2000,
10 the entire contents of which are incorporated herein by
reference.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to a method of
15 producing a biosensor protein capable of regulating a
fluorescence property of green fluorescent protein.
Furthermore, the present invention relates to
a biosensor protein (hereinafter also referred to as
a biosensor) produced by the above method and
20 a biosensor gene encoding the biosensor protein.
More specifically, the present invention relates to
a calcium-sensing biosensor protein (hereinafter also
referred to as a calcium sensor) produced by the above
method and a biosensor gene encoding the calcium-
25 sensing biosensor protein.

2. Description of the Related Art

Up to now, several biosensors have been developed

by applying the fluorescence resonance energy transfer (FRET) to green fluorescent protein (hereinafter also referred to as "GFP"). Recently, it was reported that a new type of biosensor wherein a single GFP was
5 capable of changing the fluorescence intensity without applying FRET has been developed. However, the sensitivity of the new type of biosensor was so low that it was not used in practice. To make the present invention, the present inventor has newly found that
10 the problem of the low sensitivity was caused by insufficient examination of the linking site between GFP and a functional protein employed in the biosensor.

With respect to a calcium sensor among biosensors, roughly four types of calcium sensors have been
15 developed. Hereinafter, outlines and shortcomings of these calcium sensors will be described.

1) A calcium-sensitive synthetic dye: This is a chemically synthesized dye which has sensitivity to calcium, and widely used at present. The dye is
20 loaded from the outside into a cell at the time of use. Membrane-permeable form of the dye (AM form) is easy to load into cells. However, AM form of the dye is loaded into all of the cell, so that it is difficult to introduce the dye into only a specific cell. To
25 introduce the dye into the specific cell, the dye must be injected by the use of a glass needle or the like.

2) Aequorin: This is a protein which reacts to

calcium and emits light. Aequorin is directly injected into cells at the time of use, or an aequorin-encoding gene is incorporated into a cell before use. However, aequorin requires a coenzyme to function in a cell, so that the coenzyme must be supplied to the cell. Furthermore, the light emitted therefrom is extremely weak.

3) A calcium-sensitive protein to which FRET is applied: This protein is composed of calcium-sensitive calmodulin (CaM), a partial sequence of myosin light chain kinase (which is bound to CaM), and two GFPs differing in fluorescence color. The protein utilizes the characteristics that the distance between two GFPs becomes shorter by a conformational change of CaM when calcium is bound to CaM, thereby inducing FRET, and as a result, the intensity of the fluorescence emitted from the two GFPs are altered. At the time of use, the protein is directly injected into cells, or the gene encoding the protein is incorporated into a cell. However, the change in the fluorescence intensity caused by the FRET is a little, and thus its signal is extremely weak. It is therefore hard to detect the signal by a conventional laser microscope equipped with an argon laser ($\lambda=488$ nm).

4) A calcium-sensitive protein composed of a single GFP: The calcium-sensitive protein has structure where CaM is bound between the 144th and

the 146th amino acids of the amino acid sequence
of GFP. The protein utilizes the characteristics that
the CaM protein causes a conformational change of the
GFP when calcium is bound to CaM, thereby altering the
5 intensity of the fluorescence emitted from GFP. This
protein is directly injected into a cell at the time of
use, or the gene encoding the protein is introduced
into a cell before use. However, since the calcium
sensitivity of the protein is low, so that the
10 signal/noise ratio becomes low in the cell in fact.
This fact makes it difficult to measure a signal.

BRIEF SUMMARY OF THE INVENTION

The present invention was made with the view of
overcoming the aforementioned problems. An object of
15 the present invention is to provide a method of
producing a biosensor protein capable of regulating
a fluorescence property of green fluorescent protein,
and to provide a biosensor protein produced by the
method and a biosensor gene encoding the biosensor
20 protein. Particularly, an object of the present
invention is to provide a calcium-sensing biosensor
protein produced by the method and a biosensor gene
encoding the calcium-sensing biosensor protein.

More specifically, the object of the present
25 invention is to produce a biosensor protein
(particularly a calcium-sensing biosensor protein)
whose sensitivity to a substance to be detected is high

enough to use practically and which makes it possible to measure easily. Further object of the present invention is to produce a biosensor protein (particularly a calcium-sensing biosensor protein) which is introduced into a specific cell easily and which makes it possible to measure without a specific detection device, a coenzyme, and so on.

To attain the aforementioned objects, the present invention provides the means [1] to [19] described below.

[1] A method of producing a biosensor protein capable of regulating a fluorescence property of green fluorescent protein or its derivative by modifying the structure of green fluorescent protein or its derivative, comprising the steps of:

(A) predicting a hotspot amino acid residue affecting a fluorescence property of green fluorescent protein or its derivative;

(B) producing various fusion proteins which have the structure linked with a modified fluorescent protein and one or more functional molecules,

the modified fluorescent protein being the protein obtained by cleaving amino acid sequence of green fluorescent protein or its derivative in the vicinity of the hotspot amino acid residue and modifying the structure of green fluorescent protein or its derivative, and

the functional molecules each being the molecules capable of transmitting their conformational changes to the modified fluorescent protein to cause a conformational change of the modified fluorescent protein, thereby altering the fluorescence property of the modified fluorescent protein;

(C) reacting the resultant various fusion proteins with a factor affecting the conformation of any of the functional molecules; and

(D) screening a fusion protein exhibiting a change in the fluorescence property by the reaction of the step (C), as a biosensor protein, from the various fusion proteins.

[2] The method described in [1], wherein the fluorescence property is fluorescence intensity.

[3] A biosensor protein comprising (1) and (2) below:

(1) a modified fluorescent protein which is obtained by cleaving amino acid sequence of green fluorescent protein or its derivative in the vicinity of a hotspot amino acid residue which affects the fluorescence property, and modifying the structure of the green fluorescent protein or its derivative; and

(2) one or more functional molecules which are capable of transmitting their conformational changes to the modified fluorescent protein to cause a conformational change of the modified fluorescent

protein, thereby altering the fluorescence property of the modified fluorescent protein.

[4] The biosensor protein described in [3], wherein the fluorescence property is fluorescence intensity.

[5] The biosensor protein described in [3] or [4], wherein the hotspot amino acid residue corresponds to the 148th amino acid of amino acid sequence of green fluorescent protein.

[6] The biosensor protein described in [3] or [4], wherein the hotspot amino acid residue corresponds to the 94th amino acid of amino acid sequence of green fluorescent protein.

[7] The biosensor protein described in [3] or [4], wherein the hotspot amino acid residue corresponds to the 96th amino acid of amino acid sequence of green fluorescent protein.

[8] The biosensor protein described in [3] or [4], wherein the hotspot amino acid residue corresponds to the 222nd amino acid of amino acid sequence of green fluorescent protein.

[9] A biosensor protein comprising (1) and (2) below:

(1) a modified green fluorescent protein having the following amino acid sequences (a) and (b) in this order from the N terminus:

(a) an amino acid sequence of X-th to 238th

position of green fluorescent protein; and

(b) an amino acid sequence of 1st to Y-th position of green fluorescent protein,

(where X is an arbitrary number from 148 to 150,
5 and Y is an arbitrary number from 140 to 147); and

(2) one or more functional molecules which are capable of transmitting their conformational changes to the modified green fluorescent protein to cause a conformational change of the modified green fluorescent
10 protein, thereby altering the fluorescence property of the modified green fluorescent protein.

[10] A biosensor protein comprising the following sequences (a) to (h) sequentially from the N terminus:

(a) an amino acid sequence containing methionine
15 (linker X);

(b) myosin light chain kinase protein or a partial amino acid sequence thereof;

(c) an amino acid sequence (linker Y) for linking the above sequence (b) and the following sequence (d);

20 (d) an amino acid sequence from X-th to 238th position of green fluorescent protein (where X is an arbitrary number from 148 to 150);

(e) an amino acid sequence for linking the above sequence (d) and the following sequence (f);

25 (f) an amino acid sequence from 1st to Y-th position of green fluorescent protein (where Y is an arbitrary number from 140 to 147);

(g) an amino acid sequence (linker Z) for linking the above sequence (f) and the following sequence (h); and

(h) calmodulin protein or a partial amino acid sequence thereof.

[11] A biosensor protein comprising (1) and (2) below:

(1) a modified green fluorescent protein having the following amino acid sequences (a) and (b) in this order from the N terminus:

(a) the amino acid sequence from the 149th to 238th position of green fluorescent protein, and

(b) the amino acid sequence from the 1st to 144th position of green fluorescent protein; and

(2) one or more functional molecules which are capable of transmitting their conformational changes to the modified green fluorescent protein to cause a conformational change of the modified green fluorescent protein, thereby altering the fluorescence property of the modified green fluorescent protein.

[12] A biosensor protein comprising the following sequences (a) to (h) sequentially from the N terminus:

(a) an amino acid sequence containing methionine (linker X);

(b) myosin light chain kinase protein or a partial amino acid sequence thereof;

(c) an amino acid sequence (linker Y) for linking

the above sequence (b) and the following sequence (d);

(d) the amino acid sequence from the 149th to 238th position of green fluorescent protein;

(e) an amino acid sequence for linking the above
5 sequence (d) and the following sequence (f);

(f) the amino acid sequence from the 1st to 144th position of green fluorescent protein;

(g) an amino acid sequence (linker Z) for linking
the above sequence (f) and the following sequence (h);
10 and

(h) calmodulin protein or a partial amino acid sequence thereof.

[13] The biosensor protein described in any one of [3]-[9], and [11], wherein the functional molecules
15 are calmodulin protein or a partial amino acid sequence thereof, and myosin light chain kinase protein or a partial amino acid sequence thereof.

[14] A biosensor protein comprising the following sequences (a) to (h) sequentially from the N terminus:

20 (a) Met-Gly-Thr or Met-Val-Asp (linker X);

(b) a partial amino acid sequence of myosin light chain kinase protein (Ser-Ser-Arg-Arg-Lys-Trp-Asn-Lys-Thr-Gly-His-Ala-Val-Arg-Ala-Ile-Gly-Arg-Leu-Ser-Ser);

(c) Leu-Glu (linker Y);

25 (d) the amino acid sequence from the 149th to 238th position of green fluorescent protein;

(e) Gly-Gly-Thr-Gly-Gly-Ser (linker amino acid

sequence);

(f) the amino acid sequence from the 1st to 144th position of green fluorescent protein;

(g) Gly-Thr-Arg or Thr-Arg (linker Z); and

5 (h) the amino acid sequence from the 2nd to 148th position of rat calmodulin protein.

[15] A biosensor protein comprising the following sequences (a) to (c) in this order from the N terminus:

10 (a) the amino acid sequence from the 1st to 144th position of green fluorescent protein;

(b) one or more functional molecules which are capable of transmitting their conformational changes to the modified green fluorescent protein to cause a conformational change of the modified green fluorescent protein, thereby altering the fluorescence property of the modified green fluorescent protein; and

15

(c) the amino acid sequence from the 149th to 238th position of green fluorescent protein.

[16] The biosensor protein described in [15],
20 wherein the functional molecules are calmodulin protein or a partial amino acid sequence thereof, and myosin light chain kinase protein or a partial amino acid sequence thereof.

[17] A biosensor protein comprising the following sequences (a) to (g) sequentially from the N terminus:

25

(a) the amino acid sequence from the 1st to 144th position of green fluorescent protein;

(b) an amino acid sequence (linker A) for linking the above sequence (a) and the following sequence (c);

(c) calmodulin protein or a partial amino acid sequence thereof;

5 (d) an amino acid sequence (linker B) for linking the above sequence (c) and the following sequence (e);

(e) myosin light chain kinase protein or a partial amino acid sequence thereof;

10 (f) an amino acid sequence (linker C) for linking the above sequence (e) and the following sequence (g); and

(g) the amino acid sequence from the 149th to 238th position of green fluorescent protein.

15 [18] A biosensor protein comprising the following sequences (a) to (g) sequentially from the N terminus:

(a) the amino acid sequence from the 1st to 144th position of green fluorescent protein;

(b) Gly-Thr-Arg (linker A);

20 (c) the amino acid sequence from the 2nd to 148th position of rat calmodulin protein;

(d) Gly-Thr or Gly-Thr-Gly-Ser-Gly-Gly-Gly-Ser (linker B);

25 (e) a partial amino acid sequence of myosin light chain kinase protein (Ser-Ser-Arg-Arg-Lys-Trp-Asn-Lys-Thr-Gly-His-Ala-Val-Arg-Ala-Ile-Gly-Arg-Leu-Ser-Ser);

(f) Thr-Ser (linker C); and

(g) the amino acid sequence from the 149th to

238th position of green fluorescent protein.

[19] A biosensor gene encoding the biosensor protein described in any one of [3]-[18].

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWING

5 FIG. 1A is a schematic view showing structure of GFP;

 FIG. 1B is a schematic view showing structure of a calcium sensor protein using a modified GFP;

10 FIG. 1C is a schematic view showing structure of a modified GFP; and

 FIG. 1D is a schematic view showing structure of a calcium sensor protein using a modified GFP.

 FIG. 2 shows primary structure of calcium sensor proteins (G3 and G85).

15 FIG. 3 shows a profile of the fluorescence intensity of a calcium sensor protein expressing in an HEK 293 cell, which changes in response to ATP and carbachol.

20 FIG. 4 shows a profile of the fluorescence intensity of a calcium sensor protein expressing in a skeletal muscle cell, which changes in response to carbachol and caffeine.

25 FIG. 5 is a graph showing the relationship between the calcium concentration and the fluorescence intensity.

DETAILED DESCRIPTION OF THE INVENTION

In the following, the method of producing

a biosensor protein capable of regulating the fluorescence property of green fluorescent protein of the present invention will be explained in order of the steps (A) to (E).

5 The biosensor protein of the present invention is made of a single-molecular green fluorescent protein. In the present invention, the biosensor protein may be also made of a derivative of green fluorescent protein. A derivative of green fluorescent protein means any
10 derivative of green fluorescent protein including YFP (Yellow Fluorescent Protein), CFP (Cyan Fluorescent Protein), and BFP (Blue Fluorescent Protein).

 In this specification, the biosensor protein made of green fluorescent protein will be described as
15 a representative example. Therefore, the present invention is not limited to the example.

(Method of Producing a Biosensor Protein of the Present Invention)

 (A) Step of predicting a hotspot amino acid
20 residue affecting a fluorescence property of green fluorescent protein:

 Green fluorescent protein (GFP) is a protein consisting of 238 amino acids. GFP used in the present invention is a recombinant GFP (EGFP) obtained from
25 Clontech. The EGFP contains valine (gtg) (not a constituent of original GFP) which is inserted at the back of methionine of the N terminus of original GFP.

The valine is designated as "amino acid No. 1'".

The valine (gtg) (amino acid No. 1') is omitted from the nucleotide sequence and amino acid sequence of the GFP in Sequence ID No: 1, for the sake of convenience.

5 In this text, "the n-th amino acid of green fluorescent protein" or "the amino acid of n-th position of green fluorescent protein" corresponds to the position of the 1st to 238th amino acids in the amino acid sequence described in Sequence ID No: 1.

10 In the present invention, all biosensor proteins are actually produced using the aforementioned EGFP, which has valine (amino acid No. 1') at the back of methionine of the N terminus of GFP.

15 For the sake of convenience in the present invention, the structure of GFP is shown in a schematic view of FIG. 1A. GFP has a chromophore in the center of the inside of its molecular structure. GFP can be obtained by expressing a plasmid (e.g., pEGFP-N1 (Clontech)) containing GFP-encoding cDNA.

20 In the present invention, the "hotspot amino acid residue" affecting the fluorescence property of GFP refers to an amino acid residue of GFP which acts as an indicator of a GFP modification site, in making a modified green fluorescent protein as described later.

25 The structure of GFP is modified in the vicinity of the above hotspot amino acid residue, and thereby it becomes possible to produce a desired modified green

fluorescent protein.

"Modification of GFP structure" preferably means cleaving GFP in the vicinity of the predicted hotspot amino acid residue (preferably, at any position within 5 amino acids before and behind the hotspot amino acid residue), and eliminating the appropriate number of amino acids (preferably 1 to 10 amino acids) from the cleavage site. However, the modification of GFP structure is not limited to the aforementioned modification, as long as it can produce a desired modified green fluorescent protein, and GFP may be modified by any modification process.

The desired modified green fluorescent protein (modified GFP) refers to a protein which is obtained by modifying the structure of GFP such that its fluorescence property can be reversibly changed, as described later. The phrase "fluorescence property can be reversibly changed" means that the conformation of the modified GFP reversibly changes by the presence or absence of a factor capable of causing a conformational change of the modified GFP (i.e., a biosensor-detecting factor), thereby altering the fluorescence property of the modified GFP in a reversible manner. Such reversible change of the fluorescence property refers to a change detectable by a fluorescence microscope, a laser microscope or the like, and preferably a change detectable with the naked eye. As an example of the

change in the fluorescence property, a change in the fluorescence intensity may be employed. In this case, the abovementioned change means that $\Delta F/F$ (= change amount of fluorescence intensity/initial fluorescence intensity) is preferably at least 0.1 or more, and more preferably within the range of 1 to 10.

In the present invention, the fluorescence property refers to the property such as fluorescence intensity, fluorescence wavelength, a ratio of the fluorescence intensity, absorbance, absorption wavelength, or the like. Of these, the fluorescence intensity is used herein as the fluorescence property.

The present invention is the first to find the concept of the hotspot amino acid residue. More specifically, the present inventor produced various types of biosensor proteins having a variety of modified GFPs, and thereby he found that the amino acid residue, which has a strong effect upon the fluorescence property of GFP, was present in GFP. The finding of the above amino acid residue led to the concept of the hotspot amino acid residue.

By making various biosensor proteins having a variety of modified GFPs on an experimental basis, the position of the hotspot amino acid residue can be predicted according to the performances of the obtained biosensor proteins. Also, the position of the hotspot amino acid residue can be predicted based on the

crystal structure of GFP. More specifically, there is high possibility that the hotspot amino acid residue may be an amino acid which is bound to the chromophore of GFP via hydrogen bond. Therefore, the amino acid, which is bound to the chromophore via hydrogen bond, may be predicted as the hotspot amino acid residue on the basis of the crystal structure of GFP.

In the present invention, it is predicted that the hotspot amino acid residue of GFP is an amino acid selected from the amino acids of 94th, 96th, 148th and 222nd positions of GFP. The amino acid residues of these positions are preferable as the hotspot amino acid residue, but it is not limited to these positions.

(B) Step of producing various fusion proteins which have the structure linked with a modified green fluorescent protein and one or more functional molecules,

the modified green fluorescent protein being the protein obtained by cleaving amino acid sequence of green fluorescent protein in the vicinity of the hotspot amino acid residue and modifying the structure of green fluorescent protein, and

the functional molecules each being the molecules capable of transmitting their conformational changes to the modified green fluorescent protein to cause a conformational change of the modified green fluorescent protein, thereby altering the fluorescence

property of the modified green fluorescent protein:

In the following, this step will be explained on the assumption that the hotspot amino acid residue (predicted in the step (A)) is the n-th amino acid of GFP.

In the present invention, the modified GFP preferably means those constructed by the steps below:

cleaving original GFP in the vicinity of the hotspot amino acid residue (e.g., between the n-th amino acid and the (n+1)th amino acid of GFP);

removing the amino acids in the cleavage site (e.g., amino acids from the (n-a)th to the n-th position of GFP), thereby modifying the molecular structure of GFP; and

linking the N terminus and the C terminus of original GFP with appropriate linker amino acids (e.g., Gly-Gly-Thr-Gly-Gly-Ser) in case of necessity (see FIG. 1C).

In this text, GFP (i.e., unmodified GFP) is also referred to "original GFP" in order to distinguish it from the modified GFP.

However, even if a modified GFP of a single chain is not formed by linking the N terminus and the C terminus of original GFP with the linker amino acids, two discrete GFP fragments may function in a cell, like a modified GFP of the single chain.

The linker amino acids for connecting the N

terminus and C terminus of original GFP preferably have an amino acid sequence consisting of 2 to 10 amino acid molecules. The linker amino acids more preferably have an amino acid sequence rich in glycine. Particularly
5 preferable example of the linker amino acids includes Gly-Gly-Thr-Gly-Gly-Ser. However, the linker amino acid sequence is not restricted to this example.

For example, if a modified GFP is prepared by
10 cleaving original GFP between the n -th amino acid and the $(n+1)$ th amino acid of GFP, removing the amino acids from the $(n-a)$ th to the n -th position of GFP, and linking the original N-terminus and C-terminus of GFP with the linker amino acids, the modified GFP has, in this order from the N-terminus, the $(n+1)$ th to the
15 238th amino acids, the linker amino acids, and the 1st to the $(n-a-1)$ th amino acids. In a later step, a functional molecule is linked at a newly-created N-terminus and/or C-terminus of the modified GFP (see FIG. 1D).

20 On the other hand, when the modification is performed by cleaving original GFP between the n -th amino acid and the $(n+1)$ th amino acid of GFP, removing the amino acids from the $(n-a)$ th to the n -th position of GFP, and leaving the original N terminus and C
25 terminus unlinked, the modified GFP thus obtained consists of two discrete fragments, that is, one fragment of an amino acid sequence from the 1st to the

(n-a-1)th position, and the other fragment of an amino acid sequence from the (n+1)th to the 238th position. In this case, the two discrete fragments may be linked by interposing a functional molecule between them in a later step, thereby forming a single-chain protein (see FIG. 1B). Examples of such a single-chain protein are biosensor proteins listed in Table 5 below. Even if the modified GFP does not have the structure of the single-chain protein, two discrete GFP fragments may work in a cell, like a modified GFP of the single chain.

The phrase "the vicinity of the hotspot amino acid residue (the n-th amino acid residue) at which original GFP is cleaved" refers to, for example, the range of the (n+5)th to (n-5)th position, preferably the range of the (n+2)th to (n-2)th position. Furthermore, the number of amino acids to be removed (namely, the value of a+1) is, for example, 1-10, preferably 4.

The various modified GFPs satisfying the aforementioned explanation of the modified GFP may be candidates for the biosensor protein of the present invention. However, the cleavage site in the vicinity of the hotspot amino acid residue and the number of amino acids to be removed may be appropriately set by one skilled in the art. For example, the cleavage site and the number of amino acids to be removed may be varied as shown in Tables 1 and 2.

In the present invention, the functional molecule which is linked to the modified GFP is not particularly limited, as long as it is capable of causing a conformational change itself by binding a factor acting on the functional molecule (i.e., a biosensor-detecting-factor) and it is capable of transmitting the conformational change to the modified GFP. In this text, "a conformational change" means a change of stereostructure (i.e., three-dimensional structure).

Therefore, the functional molecule used herein must be linked to the modified GFP at a position which it can transmit its conformational change to the modified GFP, thereby inducing the conformational change of the modified GFP. For this reason, the functional molecule is preferably linked to the modified GFP at a position near the modified portion thereof. More specifically, the functional molecule is preferably linked to the cleavage site of original GFP via a linker molecule.

The functional molecule may consist of a single molecule, or two or more molecules. In the case where the functional molecule consists of two molecules, the factor acting on the functional molecule (i.e., biosensor-detecting-factor) first causes a conformational change in one functional molecule of the two molecules. Next, this conformational change induces a conformational change in the other functional

molecule. Finally, each conformational change in two functional molecules can contribute to a conformational change of the modified GFP. Similarly in the case where the functional molecule consists of more than two molecules, the initial action of the factor on one functional molecule is transmitted to another functional molecule in turn, like signal transduction. Finally, each conformational change of all the functional molecules can act upon the modified GFP, causing a conformational change of the modified GFP.

As described above, in the case where the functional molecule consists of two or more molecules, conformational change each of the functional molecules can affect the structure of the modified GFP. Therefore, the effect of two or more functional molecules upon the modified GFP is larger than that of a single functional molecule. As a result, there is much possibility that the fluorescence property of the modified GFP may be drastically changed.

In the present invention, it is not always necessary that the functional molecule to be connected to the modified GFP has a complete structure as expressed in vivo. The functional molecule may have a partial structure which includes only a binding site of the factor acting on the functional molecule, as long as it can play the aforementioned role of functional molecule.

A preferable example of the functional molecule is a protein molecule. Specific examples of the functional molecule is a combination of calmodulin protein and myosin light chain kinase protein, and a combination of a CRE (Cyclic AMP Responsive Element) binding protein (CREB) and a CREB binding protein (CBP), but it is not limited to them.

For example, calmodulin protein and a part of myosin light chain kinase protein capable of binding to calmodulin (that is, M13 fragment) are employed as the functional molecules in the examples explained later. When these functional molecules are linked to the modified GFP to make a fusion protein, the fusion protein can function as a calcium sensor.

FIGS. 1B and 1D show schematic structure of the fusion proteins, wherein the functional molecules (i.e., calmodulin protein (CaM) and a part of myosin light chain kinase (M13)) are linked to the modified GFP. In addition, examples of primary structure of the fusion proteins are shown in FIG. 2.

As shown in FIG. 2, the modified GFP and the functional molecules are linked by linker molecules consisting of several amino acids. To distinguish individual linker molecules, the linker having an initiation codon (Met) at the N terminus of the fusion protein is designated as linker X; the linker connecting the N terminus side of the modified GFP

and the functional molecule (M13) is designated as linker Y; and the linker connecting the C terminus side of the modified GFP and the functional molecule (CaM) is designated as linker Z. With respect to the linker amino acids (GGTGGS) for linking the original N terminus and C terminus of GFP, the following explanation of the modified GFP can be referred to.

These linkers are not necessarily limited in length and sequence, but preferably have a length of 0 to 10 amino acids, more preferably 2 amino acids. The linker Y preferably has a sequence of leucine-glutamic acid.

Each linker is a portion for transmitting the conformational change of the functional molecule to the modified GFP. Therefore, the degree of the conformational change of the modified GFP is significantly affected depending upon length and sequence of each linker. A choice of length and sequence of each linker must be made depending upon the type of functional molecule to be linked. However, the modified GFP and the functional molecule may be directly linked without the linker.

As a matter of fact, it is necessary to produce a variety of biosensor proteins by varying length and sequence of the linker and to evaluate the performance of the biosensor proteins in experiments.

In the present invention, more specifically,

a variety of biosensor proteins are made by varying sequence of linker Y which links the modified GFP and the functional molecule (M13), and the resultant biosensor proteins are checked for their reactivity (See Tables 1 to 4).

In these biosensor proteins, when the N terminus of the modified GFP starts from the amino acid of the 149th position of original GFP, the amino acid of linker Y adjacent to the N terminal amino acid of the modified GFP is regarded as "the amino acid corresponding to the 148th position of original GFP". Furthermore, the amino acid of linker Y adjacent to "the amino acid corresponding to the 148th position of original GFP" is regarded as "the amino acid corresponding to the 147th position of original GFP".

Similarly, when the N terminus of the modified GFP starts from the amino acid of the 150th position of original GFP, the amino acid of linker Y located two amino acids away from the N terminal amino acid of the modified GFP is regarded as "the amino acid corresponding to the 148th position of original GFP".

As shown by the biosensor proteins listed on Tables 1 to 4, when "the amino acid corresponding to the 148th position of original GFP" in the amino acid sequence of linker Y has an acidic side chain (Asp of G52, Glu of G18 and G79) or a side chain having a hydroxyl group (Ser of G3 and G17, Thr of G22, Tyr of

G47), the functionally reactive biosensor may be formed. Conversely, when "the amino acid corresponding to the 148th position of original GFP" has a basic side chain (Lys of G62, Arg of G19 and G75), the reactivity of the biosensor tends to be reduced.

When "the amino acid corresponding to the 148th position of original GFP" has a hydroxyl group at the side chain, the biosensor protein further exhibits a photoisomerization property. Furthermore, "the amino acid corresponding to the 147th position of original GFP" also has an effect upon the reactivity of the biosensor protein. "The amino acid corresponding to the 147th position of original GFP" has an advantageous effect upon the reactivity of the biosensor in the order of Leu>Thr>Gly. However, when "the amino acid corresponding to the 147th position of original GFP" is Pro, it has little effect on the increase in the reactivity of Arg corresponding to the 148th position (See G19). As described above, it is possible to select an appropriate linker sequence by altering the linker sequence variously.

The biosensor protein (i.e., the fusion protein) may be formed by a known genetic engineering procedure. More specifically, first, a fusion gene is made by separately preparing each gene fragment encoding each protein to be fused (that is, a gene encoding a modified GFP and a gene encoding a functional molecule)

by polymerase chain reaction (PCR), and linking these fragments to each other. Then, the fusion protein is produced by introducing a plasmid comprising the above fusion gene into a desired cell and expressing the fusion gene.

(C) Step of reacting the resultant various fusion proteins with a factor affecting the conformation of any of the functional molecules:

As the factor affecting the conformation of a functional molecule constituting the fusion protein obtained in the step (B), it is possible to use a factor capable of causing the conformational change of the functional molecule directly by binding to the functional molecule. Alternatively, a molecule known to increase the intracellular level of the factor affecting the conformation of the functional molecule, may be used.

When the functional molecules are calmodulin protein and myosin light chain kinase, calcium ion may be used as the factor affecting the conformation of the functional molecules. Calcium ion can cause the conformational change of calmodulin by binding to calmodulin. Alternatively, it is possible to use a factor such as ATP, carbachol, caffeine, or thapsigargin, which is known to increase the calcium ion concentration in a cell expressing the fusion protein. By reacting the fusion protein with the above

factor, the function of the fusion protein as a biosensor (calcium sensor) can be confirmed.

Alternatively, when the functional molecules are a CRE binding protein (CREB) and a CREB binding protein (CBP), cyclic AMP dependent protein kinase may be used, as the factor to confirm the function as a biosensor.

Example of the method of reacting the fusion protein with the factor affecting the conformation of the functional molecule is as follows. First, cells (e.g., HEK 293 cells) producing the fusion protein of the modified GFP/functional molecules are placed in a chamber on stage of a laser microscope. Then, the factor is added to the cells. This method is preferable because the subsequent change in the fluorescence property of the fusion protein can be observed. When ATP or carbachol is used as the factor in practice, it is appropriate to suspend the factor in a perfusion solution (HBS) and add it at a concentration of 20-300 μ M.

(D) Step of screening a fusion protein exhibiting a change in the fluorescence property by the reaction of the step (C), as a biosensor protein, from the various fusion proteins:

A fusion protein exhibiting a change in the fluorescence property by reaction of the step (C) can be detected by using a fluorescence-detecting apparatus, such as a CCD camera attached to a laser

microscope or a fluorescence microscope. In this manner, the protein exhibiting a change in the fluorescence property is selected as the biosensor protein. The phrase "exhibiting a change in the fluorescence property" refers to exhibiting a change which is detectable by a fluorescence microscope, a laser microscope, or the like. More preferably, it refers to exhibiting a change sufficient to be observed by the naked eye. For example, when the change of the fluorescence property is represented by the change of the fluorescence intensity, the change amount of fluorescence ($\Delta F/F$ = change amount of fluorescence intensity/initial fluorescence intensity) is preferably 0.1 or more, more preferably falls within the range of 1 to 10.

The biosensor protein screened in the step (D) is in a state where its fluorescence property can be reversibly changed depending upon the presence or absence of the biosensor-detecting-factor.

The aforementioned biosensor protein, which is modified so that the fluorescence property of GFP can be reversibly changed, may be said to be in a critical state capable of changing the fluorescence property.

For example, when the fluorescence property is represented by the fluorescence intensity, the biosensor protein is in the critical state between a fluorescence-emitting state and

a no-fluorescence-emitting state. In the above critical state, the biosensor protein may emit no fluorescence or emit fluorescence at a low level of intensity. Alternatively, the biosensor protein in the
5 critical state may emit fluorescence at a high level of intensity.

The phrase "the biosensor protein emits no fluorescence" refers to the state where fluorescence cannot be detected by using an optical appliance.
10 The phrase "the biosensor protein emits fluorescence at a low level of intensity" refers to the state where the fluorescence intensity is so low that the biosensor protein can transfer to a level of high fluorescence intensity by the action of the
15 biosensor-detecting-factor, showing the above change amount (i.e., $\Delta F/F > 0.1$ or more). Similarly, the phrase "the biosensor protein emits fluorescence at a high level of intensity" refers to the state where the fluorescence intensity is so high that
20 the biosensor protein can transfer to a level of low fluorescence intensity by the action of the biosensor-detecting-factor, showing the above change amount (i.e., $\Delta F/F > 0.1$ or more).

When the biosensor protein is in a critical
25 state and emits fluorescence at a high level of intensity, the biosensor protein can transfer to a no-fluorescence-emitting state or weak

fluorescence-emitting state by a small conformational change thereof. Therefore, it may be said that the critical state is "the state where strong fluorescence can be barely emitted". Further, when the biosensor protein is in a critical state and emits no fluorescence, the biosensor protein can transfer to a fluorescence-emitting state by a small conformational change thereof. Therefore, it may be said that the critical state is "the state where fluorescence cannot be barely emitted". Similarly, when the biosensor protein is in a critical state and emits fluorescence at a low level of intensity, the biosensor protein can transfer to a strong fluorescence-emitting state by a small conformational change thereof. Therefore, it may be also said that the critical state is "the state where weak fluorescence can be barely emitted".

As described above, the biosensor protein of the present invention is in a critical state. This is because the biosensor protein in a critical state can sensitively respond to the presence or absence of the sensor-detecting-factor to cause a reversible change of the conformation thereof, thereby changing the fluorescence property reversibly (e.g., moving back and forth between a fluorescence-emitting state and a no-fluorescence-emitting state reversibly). Therefore, when the biosensor protein is in the above critical state, it can be said that it is in a state

where it can function as a biosensor.

(Biosensor Protein and Biosensor Gene)

The biosensor protein of the present invention refers to that screened by the aforementioned method of producing a biosensor protein. Also, the biosensor gene of the present invention refers to the gene encoding the biosensor protein of the present invention, and is used to express the biosensor protein in a cell.

The biosensor protein of the present invention is characterized by comprising (1) and (2) below:

(1) a modified GFP, which is obtained by cleaving amino acid sequence of GFP in the vicinity of a hotspot amino acid residue which affects the fluorescence property, and modifying the structure of the GFP; and

(2) one or more functional molecules, which are capable of transmitting their conformational changes to the modified GFP to cause a conformational change of the modified GFP, thereby altering the fluorescence property of the modified GFP.

The modified GFP, which is a constituent of the biosensor protein, is as defined above. The modified GFP is preferably formed by cleaving amino acid sequence of GFP in the vicinity of the hotspot amino acid residue, and eliminating some amino acids from the cleavage site, thereby modifying the molecular structure of GFP, and if necessary, linking the

original N terminus and C terminus of GFP with an appropriate amino acid sequence.

The functional molecule, which is a constituent of the biosensor protein, is also as defined above.

5 The functional molecule is a molecule capable of causing a conformational change itself by binding a factor acting upon the functional molecule (i.e., biosensor-detecting-factor), and transmitting its conformational change to the modified GFP.

10 The modified GFP and the functional molecule are not particularly limited, as long as they satisfy the aforementioned definitions.

As described above, in the biosensor protein of the present invention, the functional molecule must be
15 linked to the modified GFP at a position where the functional molecule can transmit its conformational change to the modified GFP and cause a conformational change of the modified GFP. Therefore, it is preferable that the functional molecule is linked to
20 the modified GFP in the vicinity of the modified portion of the modified GFP. More specifically, the functional molecule is preferably linked to the cleavage site of GFP via a linker molecule.

To work as the biosensor protein of the present
25 invention, the biosensor protein is required to reversibly change its fluorescence property depending on the presence or absence of the

biosensor-detecting-factor. As an example of the fluorescence property, the fluorescence intensity is used in the present invention. As described above, the biosensor protein of the present invention is in
5 a critical state between a fluorescence-emitting state and a no-fluorescence-emitting state, and can reversibly change its fluorescence property.

In the biosensor protein of the present invention, the hotspot amino acid residue is preferably the amino
10 acid residue of the 148th, 94th, 96th or 222nd position of GFP.

The biosensor protein, which is employed the amino acid of the 148th position of GFP as the hotspot amino acid residue, is characterized by comprising the
15 following (1) and (2):

(1) a modified green fluorescent protein having the following amino acid sequences (a) and (b) in this order from the N terminus:

(a) an amino acid sequence of X-th to 238th
20 position of green fluorescent protein, and

(b) an amino acid sequence of 1st to Y-th position of green fluorescent protein,

(where X is an arbitrary number from 148 to 150, and Y is an arbitrary number from 140 to 147); and

25 (2) one or more functional molecules, which are capable of transmitting their conformational changes to the modified green fluorescent protein to cause

a conformational change of the modified green fluorescent protein, thereby altering the fluorescence property of the modified green fluorescent protein.

5 More specifically, the biosensor protein, which is employed the amino acid of the 148th position of GFP as the hotspot amino acid residue, is characterized by comprising the following sequences (a) to (h) sequentially from the N terminus:

- 10 (a) an amino acid sequence containing methionine (linker X);
- (b) myosin light chain kinase protein or a partial amino acid sequence thereof;
- (c) an amino acid sequence (linker Y) for linking the above sequence (b) and the following sequence (d);
- 15 (d) an amino acid sequence from X-th to 238th position of green fluorescent protein (where X is an arbitrary number from 148 to 150);
- (e) an amino acid sequence for linking the above sequence (d) and the following sequence (f);
- 20 (f) an amino acid sequence from 1st to Y-th position of green fluorescent protein (where Y is an arbitrary number from 140 to 147);
- (g) an amino acid sequence (linker Z) for linking the above sequence (f) and the following sequence (h);
- 25 and
- (h) calmodulin protein or a partial amino acid sequence thereof.

The linker X of sequence (a) is an arbitrary amino acid sequence containing methionine, preferably an amino acid sequence consisting of 1-10 amino acids. The linker X is more preferably the sequence of
5 Met-Xaa-Xaa, wherein Xaa represents an arbitrary amino acid, and further preferably, the sequence of Met-Gly-Thr or Met-Val-Asp.

The linker Y of sequence (c) is an arbitrary amino acid sequence, preferably an amino acid sequence
10 consisting of 0-10 amino acids. More preferably, in the amino acid sequence of the linker Y, "the amino acid corresponding to the 147th position of GFP" is Leu, Thr, or Gly, and "the amino acid corresponding to the 148th position of GFP" is an amino acid having an
15 acidic side chain or a side chain with a hydroxyl group therein. Further preferably, the linker Y is the sequence of Leu-Glu.

The linker of sequence (e) is preferably an amino acid sequence consisting of 2-10 amino acids. The
20 linker of sequence (e) is more preferably an amino acid sequence rich in an amino acid (e.g., glycine) which allows a main chain of a peptide to move with a large degree of freedom, rendering the main chain flexible. Further preferably, the linker of sequence (e) is the
25 sequence of Gly-Gly-Thr-Gly-Gly-Ser.

The linker Z of sequence (g) is an arbitrary amino acid sequence, preferably an amino acid sequence

consisting of 0-10 amino acids. The linker Z is more preferably an amino acid sequence consisting of 2-3 amino acids, and further preferably the sequence of Thr-Arg.

5 In the following, Tables 1 and 2 list examples of the various biosensor proteins using the amino acid of the 148th position of GFP as the hotspot amino acid residue. Myosin light chain kinase (M13) and calmodulin (CaM) are used herein as the functional
10 molecules.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213

Table 1

| Probe No. | Linker X | M13 | Linker Y | cpEGFP | Linker Z | CaM | Fluorescence | ATP response ($\Delta F/F$) | Photoisomerization |
|-----------|----------|-----|----------|---------|----------|-----|--------------|-------------------------------|--------------------|
| G3 | MGT | O | TS | 149-144 | GTR | O | + | 0.6 | + |
| G4 | MGT | O | TS | 149-144 | GTR | CN | + | 1.6 | + |
| G6 | MGT | O | TS | 148-144 | GTR | O | - | nt | nt |
| G7 | MGT | O | TS | 147-144 | GTR | O | + | 0 | - |
| G8 | MGT | O | TS | 146-144 | GTR | O | + | 0 | - |
| G9 | MGT | O | TS | 151-144 | GTR | O | + | 0 | - |
| G10 | MGT | O | TS | 153-144 | GTR | O | * | nt | - |
| G11 | MGT | O | TS | 155-144 | GTR | O | * | nt | - |
| G12 | MGT | O | TS | 149-140 | GTR | O | + | 0 | + |
| G13 | MGT | O | TS | 148-140 | GTR | O | + | 0.2 | + |
| G14 | MGT | O | TS | 147-140 | GTR | O | * | nt | nt |
| G15 | MGT | O | TS | 146-140 | GTR | O | * | nt | nt |
| G16 | MGT | O | TS | 145-140 | GTR | O | * | nt | nt |

Table 2 (Continuation of Table 1)

| Probe No. | Linker X | M13 | Linker Y | cpEGFP | Linker Z | CaM | Fluorescence | ATP response ($\Delta F/F$) | Photoisomerization |
|-----------|----------|-----|----------|---------|----------|-----|--------------|-------------------------------|--------------------|
| G22 | MGT | O | TS | 150-144 | GTR | O | + | 0.6 | + |
| G23 | MGT | O | TS | 149-147 | GTR | CN | + | 0.2 | + |
| G24 | MGT | O | TS | 149-146 | GTR | CN | * | nt | nt |
| G25 | MGT | O | TS | 149-145 | GTR | CN | * | nt | nt |
| G27 | MGT | O | TS | 150-147 | GTR | CN | + | 1.0 | + |
| G28 | MGT | O | TS | 151-147 | GTR | CN | - | nt | nt |
| G29 | MGT | O | TS | 153-147 | GTR | CN | - | nt | nt |
| G30 | MGT | O | TS | 155-147 | GTR | CN | - | nt | nt |

The biosensor proteins listed on Tables 1 and 2 have the following amino acid sequences in this order from the N terminus:

an amino acid sequence containing methionine
5 (linker X);

a partial amino acid sequence of myosin light chain kinase protein (M13);

a linker amino acid sequence (linker Y);

a modified GFP (cpEGFP);

10 a linker amino acid sequence (linker Z); and

an amino acid sequence from the 2nd to 148th position of calmodulin protein (CaM).

The amino acid sequence of rat calmodulin used herein as a functional molecule is represented by
15 Sequence ID No: 2. Each amino acid from the 1st to 148th position of the amino acid sequence represented by Sequence ID No: 2. corresponds to the "n-th amino acid of calmodulin protein" described in this text. In the amino acid sequence represented by Sequence ID
20 No: 2, methionine of the N terminus is omitted therefrom. This is the reason why methionine of the N terminus is cleaved out by post-translational modification, after translation of genetic code of calmodulin into amino acids.

25 Also, the partial amino acid sequence (M13) of myosin light chain kinase protein, which is used as a functional molecule, is represented by Sequence

ID No: 3. The nucleotide sequence encoding the M13 fragment is obtained by modifying the nucleotide sequence encoding M13 fragment of chicken (Guerriero, V et al., Biochemistry 25, 8372-8381 (1986)).

5 In Tables 1 and 2, the names of probe No. are given to a variety of biosensor proteins listed therein. For example, the one-dimensional structure of the probe No. G3 (hereinafter, also referred to as "G3") is shown in FIG. 2. The G3 has, in this order
10 from the N terminus,

linker X (Met-Gly-Thr);

a partial amino acid sequence of myosin light chain kinase (Ser-Ser-Arg-Arg-Lys-Trp-Asn-Lys-Thr-Gly-His-Ala-Val-Arg-Ala-Ile-Gly-Arg-Leu-Ser-Ser);

15 linker Y (Thr-Ser);

the modified GFP (the amino acid sequence of the 149-238th position of GFP, the linker amino acids (Gly-Gly-Thr-Gly-Gly-Ser), and the amino acid sequence of the 1-144th position of GFP, in this order from the
20 N terminus);

linker Z (Gly-Thr-Arg); and

the amino acid sequence of the 2nd to 148th position of rat calmodulin.

25 In Tables 1 and 2, the amino acid sequences in each column of Linkers X, Y and Z are represented by single-letter notation. The symbol "O" in the column of M13 refers to the sequence of

(Ser-Ser-Arg-Arg-Lys-Trp-Asn-Lys-Thr-Gly-His-Ala-Val-Arg-Ala-Ile-Gly-Arg-Leu-Ser-Ser). The number "149-144" in the column of cpEGFP means that the modified GFP has, in this order from the N terminus, the amino acid sequence of the 149-238th position of GFP, the linker amino acids (Gly-Gly-Thr-Gly-Gly-Ser), and the amino acid sequence of the 1-144th position of GFP. The symbol "O" in the column of CaM represents the amino acid sequence of the 2nd to 148th position of original rat calmodulin, and the symbol "CN" represents the amino acid sequence of the 2nd to 148th position of calmodulin mutant CaMCN (Persechini A et al., 1997, Cell Calcium 22, 209-216). The symbol "+" in the column of Fluorescence means that a biosensor protein emits weak fluorescence before ATP is added, whereas the symbol "-" means that a biosensor protein emits no fluorescence before ATP is added. Further, the symbol "*" represents that a biosensor protein emits weak fluorescence at a high intracellular calcium concentration of about 2 mM. The value " $\Delta F/F$ " in the column of ATP response is a value (arbitrary unit) obtained by dividing " ΔF " by " F ", wherein " ΔF " is a change amount of fluorescence intensity between before and after addition of ATP and " F " is a fluorescence intensity before addition of ATP, in the cell (HEK cell) producing a biosensor protein. The symbol "nt" represents "not tested". The symbol "+" in the column

of Photoisomerization means that a biosensor protein causes photoisomerization, and the symbol "-" means that a biosensor protein causes no photoisomerization.

As shown in Tables 1 and 2, a variety of biosensor proteins are produced by varying the N terminus of cpEGFP within the amino acids of the 145-155th positions and varying the C terminus of cpEGFP within the amino acids of the 140-147th positions. The resultant biosensor proteins are checked for reactivity to ATP. As a result, the biosensor proteins having the N terminus starting from the amino acid of the 149 or 150th position, exhibit a high reactivity to ATP. The biosensor proteins having the C terminus terminating at the amino acid of 140, 144, or 147th position, exhibit a reactivity to ATP.

In some cases, it may be considered that the amino acid of the 148th position is included in linker Y. More specifically, when the N terminus of the modified GFP starts from the amino acid of the 149th position of original GFP, the amino acid of linker Y adjacent to the N terminal amino acid of the modified GFP may be regarded as "the amino acid corresponding to the 148th position of original GFP". Similarly, when the N terminus of the modified GFP starts from the amino acid of the 150th position of original GFP, the amino acid of linker Y located two amino acids away from the N terminal amino acid of the modified GFP may be regarded

as "the amino acid corresponding to the 148th position of original GFP".

5 More preferably, the biosensor protein employing the amino acid of the 148th position of GFP as the hotspot amino acid residue is characterized by comprising the following (1) and (2):

(1) a modified green fluorescent protein having the following amino acid sequences (a) and (b) in this order from the N terminus:

10 (a) the amino acid sequence from the 149th to 238th position of green fluorescent protein, and

(b) the amino acid sequence from the 1st to 144th position of green fluorescent protein; and

15 (2) one or more functional molecules, which are capable of transmitting their conformational changes to the modified green fluorescent protein to cause a conformational change of the modified green fluorescent protein, thereby altering the fluorescence property of the modified green fluorescent protein.

20 More specific example of the above more preferable biosensor protein is characterized by comprising the following sequences (a) to (h) sequentially from the N terminus:

25 (a) an amino acid sequence containing methionine (linker X);

(b) myosin light chain kinase protein or a partial amino acid sequence thereof;

(c) an amino acid sequence (linker Y) for linking the above sequence (b) and the following sequence (d);

(d) the amino acid sequence from the 149th to 238th position of green fluorescent protein;

5 (e) an amino acid sequence for linking the above sequence (d) and the following sequence (f);

(f) the amino acid sequence from the 1st to 144th position of green fluorescent protein;

(g) an amino acid sequence (linker Z) for linking
10 the above sequence (f) and the following sequence (h);
and

(h) calmodulin protein or a partial amino acid sequence thereof.

The linker X of sequence (a) is an arbitrary amino
15 acid sequence containing methionine, preferably an amino acid sequence consisting of 1-10 amino acids. The linker X is more preferably the sequence of Met-Xaa-Xaa, wherein Xaa is an arbitrary amino acid, and further preferably, the sequence of Met-Gly-Thr or
20 Met-Val-Asp.

The linker Y of sequence (c) is an arbitrary amino acid sequence, preferably an amino acid sequence consisting of 0-10 amino acids. More preferably, the linker Y is an amino acid sequence consisting of 2
25 amino acids: one is "the amino acid corresponding to the 147th position of GFP", which is Leu, Thr, or Gly; and the other is "the amino acid corresponding to the

148th position of GFP", which is an amino acid having an acidic side chain or a side chain with a hydroxyl group therein. Further preferably, the linker Y is the sequence of Leu-Glu.

5 The linker of sequence (e) is preferably an amino acid sequence consisting of 2-10 amino acids. The linker of sequence (e) is more preferably an amino acid sequence rich in an amino acid, such as glycine, which
10 allows a main chain of a peptide to move with a large degree of freedom, rendering the main chain flexible. Further preferably, the linker of sequence (e) is the sequence of Gly-Gly-Thr-Gly-Gly-Ser.

 The linker Z of sequence (g) is an arbitrary amino acid sequence, preferably an amino acid sequence
15 consisting of 0-10 amino acids. The linker Z is more preferably an amino acid sequence consisting of 2-3 amino acids, and further preferably the sequence of Thr-Arg.

 Various examples of the above more preferable
20 biosensor proteins are listed in Tables 3 and 4 in the following. Myosin light chain kinase (ML3) and calmodulin (CaM) are used herein as the functional molecules.

Table 3

| Probe No. | Linker X | M13 | Linker Y | cpEGFP | Linker Z | CaM | ATP response ($\Delta F/F$) | n | Photoisomerization |
|-----------|----------|-----|----------|---------|----------|-----|-------------------------------|----|--------------------|
| G3 | MGT | 0 | TS | 149-144 | GTR | 0 | 0.7 | 28 | +++ |
| G17 | MGT | 0 | GS | 149-144 | GTR | 0 | 0.3 | 11 | +++ |
| G18 | MGT | 0 | LE | 149-144 | GTR | 0 | 1.6 | 78 | - |
| G19 | MGT | 0 | PR | 149-144 | GTR | 0 | 0 | 12 | - |
| G41 | MGT | 0 | TI | 149-144 | GTR | 0 | 0 | 18 | - |
| G44 | MGT | 0 | TP | 149-144 | GTR | 0 | 0 | 21 | - |
| G46 | MGT | 0 | TA | 149-144 | GTR | 0 | 0 | 12 | - |
| G47 | MGT | 0 | TY | 149-144 | GTR | 0 | 0.7 | 19 | ++ |
| G49 | MGT | 0 | TQ | 149-144 | GTR | 0 | 0 | 18 | - |
| G50 | MGT | 0 | TN | 149-144 | GTR | 0 | 0 | 18 | - |
| G52 | MGT | 0 | TD | 149-144 | GTR | 0 | 0.3 | 20 | - |
| G54 | MGT | 0 | TC | 149-144 | GTR | 0 | 0.3 | 21 | - |
| G55 | MGT | 0 | TW | 149-144 | GTR | 0 | 0 | 20 | - |
| G56 | MGT | 0 | TG | 149-144 | GTR | 0 | 0 | 20 | - |
| G58 | MGT | 0 | TV | 149-144 | GTR | 0 | 0 | 23 | - |

Table 4 (Continuation of Table 3)

| Probe No. | Linker X | M13 | Linker Y | cpEGFP | Linker Z | CaM | ATP response ($\Delta F/F$) | n | Photoisomerization |
|-----------|----------|-----|----------|---------|----------|-----|-------------------------------|----|--------------------|
| G61 | MGT | O | TF | 149-144 | GTR | O | 0.5 | 22 | - |
| G62 | MGT | O | TK | 149-144 | GTR | O | 0 | 28 | - |
| G72 | MGT | O | LE | 149-144 | TR | O | 1.6 | 67 | - |
| G75 | MGT | O | TR | 149-144 | GTR | O | 0 | 29 | - |
| G76 | MGT | O | TM | 149-144 | GTR | O | 0.6 | 23 | - |
| G77 | MGT | O | TT | 149-144 | GTR | O | 0.1 | 10 | +++ |
| G79 | MGT | O | TE | 149-144 | GTR | O | 0.5 | 26 | - |
| G80 | MGT | O | TH | 149-144 | GTR | O | 0.3 | 23 | + |
| G81 | MGT | O | TL | 149-144 | GTR | O | 0.6 | 8 | - |
| G85 | MVD | O | LE | 149-144 | TR | O | 1.5 | 16 | - |
| G22 | MGT | O | TS | 150-144 | GTR | O | 0.6 | 26 | +++ |

The biosensor proteins listed on Tables 3 and 4 have, in this order from the N terminus,

an amino acid sequence containing methionine (linker X);

5 a partial amino acid sequence of myosin light chain kinase protein (M13);

a variety of linker amino acid sequence (linker Y);

a modified GFP (cpEGFP);

10 a linker amino acid sequence (linker Z); and

the amino acid sequence (CaM) of the 2nd to 148th position of calmodulin protein.

In Tables 3 and 4, the names of probe No. are given to various biosensor proteins listed therein.

15 For example, the one-dimensional structure of the probe No. G85 (hereinafter, also referred to as "G85") is shown in FIG. 2. The G85 has, in this order from the N terminus,

linker X (Met-Val-Asp);

20 a partial amino acid sequence of myosin light chain kinase (Ser-Ser-Arg-Arg-Lys-Trp-Asn-Lys-Thr-Gly-His-Ala-Val-Arg-Ala-Ile-Gly-Arg-Leu-Ser-Ser);

linker Y (Leu-Glu);

25 the modified GFP (the amino acid sequence of the 149-238th position of GFP, the linker amino acids (Gly-Gly-Thr-Gly-Gly-Ser), and the amino acid sequence of the 1-144th position of GFP, in this order from

the N terminus);

linker Z (Thr-Arg); and

the amino acid sequence of the 2nd to 148th position of rat calmodulin.

5 In Tables 3 and 4, the amino acid sequences of the linkers X, Y and Z are represented by single-letter notation. The symbol "O" in the column of M13 represents the sequence of (Ser-Ser-Arg-Arg-Lys-Trp-Asn-Lys-Thr-Gly-His-Ala-Val-Arg-Ala-Ile-Gly-Arg-Leu-Ser-Ser). The number "149-144" 10 in the column of cpEGFP means that the modified GFP has, in this order from the N terminus, the amino acid sequence from the 149-238th position of GFP, the linker amino acids (Gly-Gly-Thr-Gly-Gly-Ser), and the amino 15 acid sequence of the 1-144th position of GFP. The symbol "O" in the column of CaM represents the amino acid sequence of the 2nd to 148th position of rat-inherent calmodulin. The value " $\Delta F/F$ " in the column of ATP response is a value (arbitrary unit) 20 obtained by dividing " ΔF " by "F", wherein ΔF is a change amount of fluorescence intensity between before and after addition of ATP and "F" is a fluorescence intensity before addition of ATP, in the cell (HEK cell) producing a biosensor protein. The column "n" 25 refers to the number of cells tested. The symbol "+" in the column of Photoisomerization means that a biosensor protein causes photoisomerization, whereas

the symbol "-" means that a biosensor protein causes no photoisomerization.

In Tables 3 and 4, the reactivity of various biosensor proteins to ATP are checked by principally varying the amino acid sequence of linkers X, Y, and Z. Biosensor proteins having the sequence of Leu-Glu as linker Y show the highest reactivity. In both cases where the sequence of linker Z is (Gly-Thr-Arg) and (Thr-Arg), there is no significant difference between them with respect to the reactivity to ATP. Furthermore, in both cases where the sequence of linker X is (Met-Gly-Thr) and (Met-Val-Asp), there is no significant difference between them with respect to the reactivity to ATP. Although the total number of the amino acids of cpeGFP (150-144) in the biosensor protein G22 is smaller than that of cpeGFP (149-144) in G3 by one, there is no significant difference between them in the reactivity.

As shown in the results of Tables 3 and 4, the biosensor proteins G18, G72, and G85 exhibit the very high reactivity, and show the value 1.5-1.6 in the change amount ($\Delta F/F$) of fluorescence to ATP. All of the biosensor proteins G18, G72, and G85 have the modified GFPs having the N terminus starting from the 149th amino acid of original GFP, and have glutamic acid (Glu) as "the amino acid corresponding to the 148th position of GFP" in linker Y. In addition,

the results show that the biosensor protein G22 exhibits a relatively low reactivity ($\Delta F/F=0.6$), but has a sufficient sensitivity to calcium. The biosensor protein G22 has the modified GFP having the N terminus starting from the 150th amino acid of original GFP, and has threonine as "the amino acid corresponding to the 148th position of GFP" in linker Y.

From the results of Tables 3 and 4, the following facts are also demonstrated. When "the amino acid corresponding to the 148th position of GFP" in linker Y has a side chain with a hydroxyl group, such biosensor protein causes photoisomerization. This suggests that the amino acid of linker Y adjacent to the modified GFP (i.e., "the amino acid corresponding to the 148th position of GFP") is closely related to the fluorescence intensity of the biosensor protein.

Other than the biosensor proteins described in Tables 1-4, another examples of biosensor proteins are listed on Table 5. The biosensor proteins listed on Table 5 are prepared by cleaving GFP in the vicinity of the amino acid residue of the 148th position of GFP (which is the hotspot amino acid residue), and linking the functional molecules between the cleaved sites.

Table 5

| Probe No. | N-terminal of modified GFP | Linker A | CaM | Linker B | M13 | Linker C | C-terminal sequence of modified GFP | Fluorescence | ATP response ($\Delta F/F$) | Photoisomerization |
|-----------|----------------------------|----------|-----|----------|-----|----------|-------------------------------------|--------------|-------------------------------|--------------------|
| A1 | 1-144 | GTR | O | GT | O | TS | 149-238 | + | 0.3* | + |
| A2 | 1-144 | GTR | O | GTSGGGG | O | TS | 149-238 | + | 0.5 | + |

The biosensor proteins listed on Table 5 have the following sequences in this order from the N terminus:

the amino acid sequence of the 1st to 144th position of green fluorescent protein;

5 linker A;

a partial amino acid sequence (M13) of myosin light chain kinase;

linker B;

10 the amino acid sequence (CaM) of the 2nd to 148th position of rat calmodulin;

linker C;

the amino acid sequence of the 149 to 238th position of GFP.

15 In Table 5, the amino acid sequences of the linkers A, B and C are represented by single-letter notation. The symbol "O" in the column of M13 represents the sequence of (Ser-Ser-Arg-Arg-Lys-Trp-Asn-Lys-Thr-Gly-His-Ala-Val-Arg-Ala-Ile-Gly-Arg-Leu-Ser-Ser). The symbol "O" in
20 the column of CaM represents the amino acid sequence of the 2nd to 148th position of rat-inherent calmodulin. The symbol "+" in the column of Fluorescence means that a biosensor protein emits weak fluorescence before addition of ATP. The value " $\Delta F/F$ " in the column of
25 ATP response is a value (arbitrary unit) obtained by dividing " ΔF " by "F", wherein " ΔF " is a change amount of fluorescence intensity between before and after

addition of ATP and "F" is a fluorescence intensity before addition of ATP, in the cell (HEK cell) producing a biosensor protein. The symbol "*" in the column of ATP response represents that the biosensor protein responds to carbachol in the case of using carbachol instead of ATP. The symbol "+" in the column of Photoisomerization means that a biosensor protein causes photoisomerization.

Table 5 shows the ATP reactivity of a biosensor group having the original N terminus and original C terminus of GFP. The biosensor group reacts to ATP, but the reactivity is low. The biosensor protein A2 having a longer linker B shows higher reactivity than A1. The reactivity of this biosensor group will be improved by examining the various sequences with respect to linkers A, B and C, and a biosensor protein having higher reactivity will be produced.

The results of Tables 1 to 5 reveal that GFP has the amino acid residue of the 148th position as the hotspot amino acid residue which has a strong effect upon the fluorescence intensity of GFP. This makes it possible to control the fluorescence intensity of GFP efficiently by way of the hotspot amino acid residue. Similarly, it is predicted from the crystalline structure of GFP that a biosensor protein capable of controlling the fluorescence intensity of GFP can be constructed, if the 94th, 96th, or 222nd amino acid

residue is employed as the hotspot amino acid residue, as well as the 148th amino acid residue.

Examples of the biosensor proteins of the present invention include the biosensor proteins having
5 calmodulin protein (or a part thereof) and myosin light chain kinase (or a part thereof) as the functional molecules, as described above. However, the biosensor proteins of the present invention are not limited to them. In other words, the biosensor proteins of the
10 present invention are not limited to those working as the aforementioned calcium sensor.

Particularly preferable example of the calcium sensor protein of the present invention is characterized by comprising the following sequences (a)
15 to (h), sequentially from the N terminus,

- (a) Met-Gly-Thr or Met-Val-Asp (linker X);
- (b) a partial amino acid sequence of myosin light chain kinase protein (Ser-Ser-Arg-Arg-Lys-Trp-Asn-Lys-Thr-Gly-His-Ala-Val-Arg-Ala-Ile-Gly-Arg-Leu-Ser-Ser);
- 20 (c) Leu-Glu (linker Y);
- (d) the amino acid sequence of the 149th to 238th position of green fluorescent protein;
- (e) Gly-Gly-Thr-Gly-Gly-Ser;
- (f) the amino acid sequence of the 1st to 144th
25 position of green fluorescent protein;
- (g) Gly-Thr-Arg or Thr-Arg (linker Z); and
- (h) the amino acid sequence from the 2nd to 148th

position of rat calmodulin protein.

The calcium sensor prepared in the present invention can be used when calcium concentration is measured inside or outside cells.

5 For example, calcium concentration can be measured by previously preparing a calcium sensor protein of the present invention by introducing a gene encoding the calcium sensor protein into *E. coli* or the like, and then mixing the resultant calcium sensor protein with
10 a specimen. Further, the intracellular calcium concentration may be measured by injecting directly the calcium sensor protein produced by using *E. coli* or the like into a desired cell to be measured for calcium concentration. Alternatively, the intracellular
15 calcium concentration may be measured by introducing the calcium sensor gene into the desired cell to be measured for calcium concentration, and expressing the gene in the cell.

 The measurement of calcium concentration can be
20 performed by irradiating light at a specific wavelength (e.g., excitation light at 488 nm) to a calcium sensor protein and detecting the property of the fluorescence emitted from the calcium sensor protein by means of an optical appliance (e.g., laser microscope). As regards
25 the calcium sensor protein used for measurement, it is necessary to previously check the relationship between a known calcium concentration and the fluorescence

property of the calcium sensor protein under the known calcium concentration. More specifically, for example, it is necessary to make a large amount of the calcium sensor protein by using *E. coli*, and previously measure the property of fluorescence wavelength and the change of fluorescence intensity in response to various calcium concentration by fluorescent spectrophotometer (See FIG. 5). The biosensor protein of the present invention has almost the same property of fluorescence wavelength as that of EGFP (Clontech). Specifically, the biosensor protein of the present invention has a maximum excitation wavelength of 489 nm and a maximum emission wavelength of 509 nm.

In the present invention, the biosensor gene refers to a gene encoding the biosensor protein of the present invention. The biosensor gene is prepared, in the form of a fusion gene, by making each gene fragment encoding individual constitutional parts (e.g., parts of a modified GFP and a functional molecule) of a biosensor protein by means of PCR, and linking the resultant each gene fragment. The biosensor gene thus prepared is introduced into an appropriate vector (pEGFP-N1, Clontech), and then the recombinant plasmid is further introduced into a cell. In this manner, the desired biosensor protein can be produced in an arbitrary cell.

(Confirmation of the Performance of the Biosensor Protein of the present invention)

The biosensor protein of the present invention is required checking experimentally for the performance as a sensor in advance.

The calcium sensor which was actually developed in the present invention was checked for the performance, by introducing the gene of the calcium sensor into cultured human embryonic kidney (HEK) 293 cells or mouse skeletal muscle primary cultured cells.

The calcium sensor of probe No. G85 is made by expressing the gene of the calcium sensor in the HEK 293 cells, and then a factor known to increase the calcium ion concentration in cells, such as ATP (0.1 mM) or carbachol (CCH) (0.1 mM), is added to the HEK 293 cells. The reaction profile of the calcium sensor G85 is shown in FIG. 3. The nucleotide sequence of the G85 is represented by Sequence ID No: 4

As shown in FIG. 3, the change amount ($\Delta F/F$) of fluorescence in response to ATP in the HEK 293 cells comes up to about 1.6. Such change amount can be captured by both fluorescence microscopy and naked eye. The change amount ($\Delta F/F$) of fluorescence in response to carbachol reaches about 2.2. Furthermore, it is observed that the fluorescence intensity of the calcium sensor reaches a maximum, when ionomycin (0.01 mM) and a calcium ion (2 mM) are added to increase the calcium

concentration in the cells. Conversely, it is observed that the fluorescence intensity of the calcium sensor reaches a minimum, when the calcium concentration in the cells is approximated to zero by adding ionomycin (0.01 mM) and EDTA (Ethylenediamine-N,N',N' tetraacetic acid disodium salt) (10 mM), without adding a calcium ion.

Similarly, a factor such as carbachol (0.1 mM) or caffeine (10 mM), which is known to increase the intracellular calcium ion concentration, is added to the skeletal muscle primary cultured cells expressing a calcium sensor (G85). The profile of the reaction is shown in FIG. 4. The change amount ($\Delta F/F$) of fluorescence in response to carbachol comes up to about 2.5, and the change amount ($\Delta F/F$) of fluorescence in response to caffeine reaches about 1.1, which is similar to the result of FIG. 3.

Furthermore, the relationship between the calcium concentration and the fluorescence amount of the calcium sensor (G85) is shown in the graph of FIG. 5. The G85 is prepared by using E. coli. As shown in FIG. 5, the calcium sensor protein of the present invention exhibits extremely high calcium sensitivity. This sensitivity is about 30 times as high as that of a calcium-sensitive protein consisting of a single GFP described in the column of "BACKGROUND OF THE INVENTION".

As described above, it is demonstrated that the calcium sensor of the present invention has highly efficient performance as a sensor.

(Examples)

5 Method of Preparing Calcium Sensor and Method of using the Calcium Sensor

Construction of pN1-G3 encoding a probe No. G3

1. Smooth Muscle Myosin Light Chain Kinase M13

20 The fragment M13 (Sequence ID No: 3) of smooth muscle myosin light chain kinase was prepared using the following synthetic primers which have been synthesized by Katayama Chemical. More specifically, PCR was performed using the following two primers as templates and primers, thereby synthesizing the fragment M13 of
15 smooth muscle myosin light chain kinase.

 smMLCKM13-1 primer:

 GCGCTAGCCGCCACCATGGGTACCTCATCACGTCGTAAGTGGGAATAAGAC
AGGTCACGCAGTCAGA (Sequence ID No: 5)

 smMLCKM13-2 primer:

20 GGCGCGGCGGCTCAACTAGTTGAGCTCAGCCGACCTATAGCTCTGACTGC
GTGACCTGTCTT (Sequence ID No: 6)

 The reaction mixture obtained by PCR was subjected to agarose gel electrophoresis. A target PCR product was recovered from the gel, and digested with NheI and
25 SpeI restriction enzymes, to obtain a PCR fragment 1.

2. Modified GFP (i.e., Circularly Permutated Enhanced GFP: cpEGFP)

PCR was performed using the following primers and a plasmid pEGFP-N1 (available from Clontech) containing GFP-encoding cDNA as a template.

EGFP-31 primer:

5 GGACGCGTACTAGTAACGTCTATATCATGGCCGAC (Sequence ID No: 7)

EGFP-20 primer:

CCGGTACCGCCCTTGTACAGCTCGTCCATGCC (Sequence ID No: 8)

10 The reaction mixture obtained by PCR was subjected to agarose gel electrophoresis. A target PCR product was recovered from the gel, and digested with SpeI and KpnI restriction enzymes, to obtain a PCR fragment 2.

15 Similarly, PCR was performed using the following primers and a plasmid pEGFP-N1 as a template.

EGFP-21 primer:

GCGGTACCGGAGGGAGCATGGTGAGCAAGGGCGAGGAG (Sequence ID No: 9)

EGFP-30 primer:

20 GGACGCGTCCCGTTGTACTCCAGCTTGTGCCC (Sequence ID No: 10)

25 The reaction mixture obtained by PCR was subjected to agarose gel electrophoresis. A target PCR product was recovered from the gel, and digested with KpnI and MluI restriction enzymes, to obtain a PCR fragment 3.

3. Calmodulin (CaM)

PCR was performed using the following primers to

rat calmodulin and a plasmid rCaM encoding a rat calmodulin cDNA (Sequence ID No: 2; Accession No. M19312 (Mori M, 2000, Biochemistry 39, 1316-1323)) as a template.

5 rCaM-2 primer: GGACGCGTGACCAACTGACTGAAGAGCAG
(Sequence ID No: 11)

 rCaM-10 primer: GCGCGGCCGCTCACTTCGCTGTCATCATTGTAC
(Sequence ID No: 12)

10 The reaction mixture obtained by PCR was subjected to agarose gel electrophoresis. A target PCR product was recovered from the gel, and digested with MluI and NotI restriction enzymes, to obtain a PCR fragment 4.

15 The four fragments obtained were linked stepwise with DNA ligase, and the ligated fragment was finally introduced into a pEGFP-N1 vector which was previously digested with NheI and NotI restriction enzymes.

 More specifically, the above four types of PCR fragments were prepared as described below.

20 The primers used in each PCR were synthesized primers (Sawady and Katayama Chemical).

 The conditions in each PCR were as follows.

| | | |
|----|---|------------------|
| 25 | Template DNA | 1 μ g |
| | Two primers | 10 pmol for each |
| | Reaction buffer (Stratagene) | 5 μ L |
| | Pfu enzyme (Stratagene) | 2 units |
| | 2.5 mM dNTP (Takara) | 4 μ L |
| | Water was added to adjust a total volume to | |

50 μ L.

The reaction temperature in each PCR was as follows.

- 1) 94°C, 2 minutes, 1 cycle
- 5 2) 94°C, 30 seconds
 55°C, 30 seconds
 72°C, 1 minute, 30 cycles
- 3) 72°C, 2 minutes, 1 cycle

10 The recovery of DNA from gel was performed by
using DNA Fragment Purification Kit MagExtractor
(Toyobo) in accordance with the manual attached to the
kit.

15 The agarose gel electrophoresis was performed
using 1% or 2% agarose gel (Agarose LE, Nakarai Tesque)
and TAE buffer (4.98g/L Tris base (Nakarai Tesque),
1.142 mL/L glacial acetic acid (Nakarai Tesque), and
2 mL/L of 0.5 M EDTA (pH 8) (Dojin)).

20 The cleavage of DNA by restriction enzymes was
performed using any of NheI, SpeI, MluI, and NotI
(available from NEB) and buffer attached to the
enzymes. Specifically, the enzymes (30 units for
each), the buffer (10 μ L), and water were added to
1-2 μ g of DNA, to adjust the total amount of the
reaction mixture to 100 μ L, and then the resultant
25 reaction mixture was reacted for 1-3 hours at 37°C.
Thereafter, DNA was recovered by using DNA Fragment
Purification Kit MagExtractor (Toyobo) in accordance

with the manual attached to the kit.

The ligation reaction was performed using a DNA Ligation kit (Takara) in accordance with the manual attached to the kit.

5 E. coli HB101 (Takara) was used as a competent cell, and transformed by a calcium chloride method (Molecular Cloning A Laboratory Manual, 2nd Edition, by J.Sambrook, E.F.Fritsch, T.Maniatis, Cold Spring Harbor Laboratory Press, 1989).

10 The recovery of the plasmid from E. coli was performed by using QIAprep Spin Miniprep kit (Qiagen) in accordance with the manual attached to the kit.

Introduction of plasmid to cells:

15 cDNA (that is, the plasmid recovered from E. coli) was introduced into HEK 293 cells (American Type Culture Collection) placed on a cover glass of 5 × 10 mm, by the use of a Superfect transfection reagent (Qiagen) in accordance with the manual attached thereto. The HEK 293 cells used herein were previously
20 incubated in Dulbecco's modified Eagle medium (Gibco BRL) containing 10% fetal bovine serum, penicillin (30 units/mL), and streptomycin (30 mg/mL).

More specifically, the cDNA (that is, the plasmid recovered from E. coli) was introduced as follows:
25 First, for HEK cells cultured in 3 cm-dish, 12.5 μ L of Superfect transfection reagent and 80 μ L of serum-free Dulbecco's modified Eagle medium were added to 3 μ g of

the cDNA (3 μ L). The resultant reaction mixture was allowed to stand still at room temperature for 10 minutes. Thereafter, to the reaction mixture, 800 μ L of serum-containing Dulbecco's modified Eagle medium was added. The resultant reaction mixture was added to the HEK cells which was previously washed once with a phosphate buffer, and then the HEK cells were incubated for 2 hours at 37°C. The incubated HEK cells were washed once with the phosphate buffer.

Thereafter, the Dulbecco's modified Eagle medium containing 10% fetal bovine serum was added to the HEK cells in an appropriate amount (2 mL per 3cm-dish), and the cells were incubated at 28°C for 2-4 days. The obtained cells were used in experiments.

Measurement:

The cover glass having the obtained cultured cells attached thereto was placed in a chamber on a stage of microscope, and refluxed with HBS solution (107 mM NaCl, 6 mM KCl, 1.2 mM MgSO₄, 2 mM CaCl₂, 1.2 mM KH₂PO₄, 11.5 mM glucose, and 20 mM HEPES pH 7.4 (purchased from Nakarai Tesque, Katayama Chemical, HEPES: 2-[4-(2-Hydroxyethyl)-1-piperazinyl]ethanesulfonic acid, Dojin)). 100 μ M of adenosine 3'-phosphate (ATP, Sigma) or 100 μ M of carbachol (Sigma) was dissolved in HBS, and the resultant ATP or carbachol solution was supplied to the cells by reflux. Using a Leica TCS-NT laser

microscope, the cells were excited at a wavelength of 488 nm, and the emitted fluorescence was recorded at a wavelength of 525 nm.

(Effects of the Invention)

5 As explained in the foregoing, the method of producing a biosensor protein of the present invention can be applied to the development of various types of biosensors. Also, it is expected that a high-performance biosensor can be developed in a short
10 period by using the method of producing a biosensor protein of the present invention.

The change amount in the signal intensity of the calcium sensor produced by using the method is sufficiently large to be observed by the naked eye.
15 The calcium sensitivity of the calcium sensor is improved to 30 times as large as that of a conventional calcium sensor. Furthermore, since the calcium sensor is a protein sensor, the sensor can be readily expressed in a cell by introducing the sensor-encoding
20 gene into the cell. Furthermore, the present invention has the following advantages: A specific device and substance are not required for measurement, and a detector using an argon laser can be used. Also, a coenzyme is not required for measurement.

25 In addition, the calcium sensor developed herein can be incorporated into a instrument for measuring a calcium concentration, by the use of the sensor

protein itself. Moreover, the animal, plant, tissue,
organ, cell, or the like, to which the gene encoding
a biosensor protein of the present invention is
introduced, can be used in developing pharmaceutical
5 and agricultural products.

Additional advantages and modifications will
readily occur to those skilled in the art. Therefore,
the invention in its broader aspects is not limited to
the specific details and representative embodiments
10 shown and described herein. Accordingly, various
modifications may be made without departing from the
spirit or scope of the general inventive concept as
defined by the appended claims and their equivalents.